

PERENNIAL

PROPAGATION

Buddleia Nanho Series

Buddleia davidii



TRAY & ROOTING

Form	Unrooted Cuttings (URC)
Rooting Hormone	Yes
Tray Size(s)	50, 105
Plants Per Cell	1
Wks. From Stick To Transplant	4-5
Pinch?	Yes
Days from stick to pinch	14-21

FERTILITY

Fertilize Through Mist?	No
Fertilizer Rate	100-150 ppm N (20-6-22 or 22-5-16 [15-0-15 under low-light conditions])
EC Range	Moderate: (2.0-3.5 mS via SME)
pH Range	5.8-6.2

TEMPERATURE

Average Soil Temperature	18-22°C)
Average Air Temperature	18-22°C

IRRIGATION

Irrigation Frequency	Level 3: Keep media moderately moist with mild dry cycles
----------------------	--

LIGHT

Supplemental	No
Benefits From Shade?	No

POTENTIAL DISEASES

Botrytis, downy mildew

POTENTIAL PESTS

aphids, whitefly

PGR SUGGESTIONS

- Optional 300-500 ppm Florel (3.9% Ethephon)
- Optional 3-5 ppm Sumagic (0.055% Uniconazole)

CULTIVAR LIST

Blue, Purple



'Purple'

TIPS

- Cuttings are sensitive to drying out; tip burn is a tell-tale sign of this issue.

WE'RE HERE TO HELP YOU GROW

Robbin den Herder- Product Manager
R.denHerder@DummenOrange.com
+31 174 530 100
[DummenOrange.com](https://www.DummenOrange.com)

PERENNIAL
FINISHING
Buddleia Nanho Series

Buddleia davidii



VERNALIZATION

Vernalization Required?	No
Vernalization Duration Minimum	N/A
Vernalization Temperature	N/A

CONTAINER SIZE	PLANTS PER POT	TRANSPLANT TO FINISH (WEEKS)
----------------	----------------	------------------------------

16 cm	1	10-12
22 cm	2	10-12
30 cm	3	10-12

FERTILITY

Fertilizer Rate 150-200 ppm N (20-6-22 or 22-5-16 [15-0-15 under low-light conditions])

EC Range **High: (3.5-5 mS via SME)**

pH Range **5.8-6.3**

TEMPERATURE

Day Temperature **18-24°C**

Night Temperature **13-18°C**

IRRIGATION

Irrigation Frequency **Level 4: Keep media moderately moist at all times**

LIGHT

Photoperiodic requirements **Long day: 12+ hours daily**

Supplemental? **Daylength Extension**

Benefits From Shade? **No**

PINCHING

Finishing Pinch? **Optional**

Transplant to pinch (days) **14-21**

POTENTIAL DISEASES

Botrytis, downy mildew

POTENTIAL PESTS

aphids, whitefly

PGR SUGGESTIONS

- Bonzi (0.4% Paclobutrazol) 5-40ppm spray
- 5-8 ppm Sumagic (0.055% Uniconazole)
- 2500 ppm B-Nine (85% Daminozide) as needed



'Pink'

TIPS

- Final pinch 4 weeks after transplanting. Heavy PGR applications are required.
- Full sun (preferably outdoor production) and drip irrigation is the best system for finishing.
- The large and heavy root system requires additional irrigation. Make sure Buddleia is not allowed to dry- out completely.
- Avoid fertilizing during winter months when the plant isn't actively growing. This will cause a buildup of soluble salts in the media and increase risk of root burn.

WE'RE HERE TO HELP YOU GROW

Robbin den Herder- Product Manager
R.denHerder@DummenOrange.com
+31 174 530 100
DummenOrange.com

PERENNIAL

FEATURES AND BENEFITS

Buddleia Nanho Series

Buddleia davidii



KEY INFORMATION

First Year Flowering?	Yes
Bloom Time	Early Summer to Late Summer
Exposure	Full Sun
Hardiness Zone	5-11
Height	122-152 cm
Width	152 cm

DESCRIPTION

Buddleia has regained popularity with new very compact breeding and bright colors. The classic Buddleia architecture is still widely appreciated for its quick and dramatic impact in the landscape.

CONSUMER APPEAL

- As the common name implies, Buddleia are the ultimate butterfly attractors but they also support bee species and hummingbirds as well. Due to its shrubby nature, Butterfly Bush is generally avoided by deer and rabbits.

GARDEN ATTRIBUTES

- 'Nanho' is a vigorous series with a broad arching habit giving it exceptional impact in the landscape.
- Although unusual and somewhat unorthodox, Buddleia blooms have a wholesome "fresh-baked-bread" fragrance to their flowers.
- Remarkably drought tolerant once established, these sub-shrubs aren't picky when it comes to soil conditions; happy with dry or moist. If planting in clay soil, amend the soil first with compost or other organic matter to improve the drainage.
- In northern climates, Buddleia is classified as a "sub-shrub" meaning that it dies back to the ground during the winter and reestablishes itself from the root system the following year. To ensure a good growing habit, remove the previous year's woody stem

NATIVITY

China

COMPANION PLANTS

Ceratostigma, Glechoma, Lamium, Lamiastrum, Mazus, Verbena, Vinca, Lysimachia, Lithodora, Iberis, Hypericum, Hedera, Geranium

COMMON NAME(S)

butterfly bush, summer lilac



'Nanho Purple'

WE'RE HERE TO HELP YOU GROW

Robbin den Herder- Product Manager

R.denHerder@DummenOrange.com

+31 174 530 100

DummenOrange.com