



SINGLE



DECO



SPECIALTY



OUTDOOR

# KALANCHOE

## Culture Guide

**Content:**

- Ten commandments of kalanchoe growing
- Temperature
- Humidity
- Micro climate
- Substrate
- Long days
- Short days
- Rooting
- Irrigation
- Fertilization
- EC schedule
- CO<sup>2</sup>
- Lighting
- Light
- Shading
- Growth retardents (in general)
- Growth retardents (Daminozide)
- Schedule for Daminozide treatment (basic)
- Spraying methods
- Long Day period
- Short Day period (induction period)
- Daminozide/Bonzi
- Crop protection
- Biological control
- Mother stock culture
- Culture decoratives
- Outdoor Culture (Calor/Deck'd Out)
- Culture in high temperatures

## **TEN COMMANDMENTS OF KALANCHOE GROWING**

1. Keep a steady balance in the culture in every aspect, avoid as possible any shocks.
2. In the Long Days, keep the culture as dry as possible for the first 8-14 days. No misting.
3. Growth regulators in the Long Day period, if applied a misty treatment is the right way.
4. At least 14 hours of darkness is needed for induction, for a period of at least 7 weeks.
5. If you can read a newspaper under blackout, it is not dark enough.
6. Short Day interruption is impossible, the moonlight can already disturb induction.
7. In the Short Day period growth regulators are at least essential in the induction period.
8. Stretching in Kalanchoe is mainly caused by humidity.
9. Avoid water overhead in the Short Day, never use cold water (under 10°C/50°F)
10. Don't use oil based liquid chemicals, unless advised by experts.

## **TEMPERATURE**

1. Ideal temperature is 20 °C (68 °F).
2. Every degree lower than 20 °C (68 °F) gives a week delay.
3. Every degree lower than 20 °C (68 °F) means one week less in keeping quality.
4. Lower temperatures increase the flower size (Think about items 2 and 3).
5. Daytime Temperatures over 32 °C (89 °F) cause extra stretching.
6. Nighttime temperatures of 23 °C (73 °F) or higher cause disturbances in or no induction.
7. Temperatures under 16 °C (60 °F) cause disturbances in or no induction.
8. Negative Dif only gives growth reduction if it is inside 3 degrees. (Think about items 2 and 3).

## **HUMIDITY**

1. Ideal is around 85% (70-90%).
2. More than 25% difference is the main cause for stretching.
3. Lower than 40% (dry climate) creates problems, weaker plant.
4. Higher than 95% creates problems, softer plant.
5. Condensating of moisture gives a higher risk of fungi infections.

## **MICRO-CLIMATE**

1. Micro-climate is more important than greenhouse climate.
2. Pad and Fan system in greenhouses take away the micro-climate on plant level.
3. Irrigation mat improves micro-climate on table and container systems.

## **SOIL**

1. Must be airy.
2. Must have draining capacity.
3. Must have a storing capacity.
4. pH 5.5 – 6.0.
5. Choice of soil depends on growing system.
6. Quality soil needed for quality roots

## **LONG DAYS**

1. Rooting period as dry as possible.
2. Rooting hormones not necessary.
3. No misting.
4. Covering for rooting with poly not necessary.
5. Covering for protection against radiation possible.
6. Ideal temperature 21-22 degrees °C (70 – 74 degrees °F)
7. Pinching not necessary.
8. Mist treatment of growth retardant necessary on 2<sup>nd</sup>/early 3<sup>rd</sup> week.  
Purpose: form of chemical pinching (slow down headshoot, stimulating sideshoots). Growth retardant 4<sup>th</sup> week, controls of head size.
9. Not more than 18 hours of light.
10. Natural Long Day: end march – half september.

## **SHORT DAYS**

1. 14 hours of darkness.
2. Natural Short Day: half september – half march.
3. 3<sup>d</sup> – 6<sup>th</sup> week: induction period.
4. Induction period: important growth retarding moment.
5. Length of short day: minimum 6 weeks.
6. Preference: short day till flowering/sales.
7. Light level under covering: < 10 lux is perfect.
8. Short day interruption not possible.
9. Interruption light: streetlanterns, neighboring greenhouses, the moon.
10. Not more than 18 hours of light.

## **ROOTING**

1. Fill the pots with soil.
2. Drench to pots, make them wet.
3. Wait for half a day before sticking the cuttings.
4. No covering with plastic or tissue necessary (choice).
5. No water for at least 4 - 7 days.
6. If the soil dries out, just give a short shower.
7. After 12 days, start normal irrigation and feeding.


## **IRRIGATION**

1. No misting in the Long Day period.
2. Keep the culture moist, not wet.
3. High frequency (every day a little) is better than low frequency (once a week a lot).
4. High temperatures: cooling off shower (specificly high noon).
5. Too dry: anthocyanin (red coloring) on underside leaves.
6. Too wet: drowning of roots.

## FERTILIZATION

1. Ideal pH 5.8 (range 5.5 – 6.1)
2. Ideal EC Long Days: 1.0.
3. Maximum EC allowed: 1.5
4. EC Short Days – induction: 1.0 -> 1.5.
5. EC Short Days – last weeks: 1.5 -> 1.0.
6. Hot conditions (over 25 °C daytime/ 76 °F): lower EC required.
7. Important elements: Calcium (Ca), Potassium (K), Phosphorus (P), Nitrogen (N) and Magnesium (Mg).

## EC SCHEDULE CULTURE (IN POT)



- a) In normal conditions decreasing the EC from the first period of the Short Day to the last period can take a couple of weeks.
- b) When the temperature is high, over 28°C (82 °F) outside, decreasing the temperature has to be quicker.

## CO<sub>2</sub>

1. CO<sub>2</sub> is essential for branching
2. Over 1100 ppm causes problems.
3. Under 400 ppm (natural level) means lack of growth.
4. Kalanchoe is a CAM (Crassulacean Acid Metabolism), which means it can in CO<sub>2</sub> at night as well.
5. Apply CO<sub>2</sub> mainly at night in the first two weeks (rooting) and the last weeks (flowering).
6. Apply CO<sub>2</sub> mainly in the day in the last period of Long Days and the first period of Short Days (including induction).

## LIGHTING

1. Assimilation lights = grow light.
2. Cyclic light is not grow light: 10 minutes per half an hour.
3. Night break most ideal.
4. Start 2-3 hours after sunset, ending 2-3 hours before sunrise.
5. Minimum 15 Watt per m<sup>2</sup> (100 lux 3).
6. Distance light – plant level: 2-2.5 meter.
7. Daytime support light: maximum 10 hours.

## LIGHT

1. Critical daylength: 11,5 hours.
2. Daylength: minimum 13 hours, maximum 18 hours.
3. Natural light period: end march – half september.
4. Too much light: anthocyanin (red coloring) on leaves.

## SHADING

1. Summer: starting at 55.000 lux.
2. Spring: starting at 45.000 lux.
3. Calculate shading with starting point high noon (12 o'clock).

## GROWTH RETARDANTS (IN GENERAL)

1. Some similar growth retardants are different on level of active ingredient. F.i. Bounty in Japan has approximately 40 times more active ingredient than Bonzi.
2. Some growth retardants compete with the hormone responsible for making anthocyanin in the plant. Discoloring (to lighter colors) can be the result of that.
3. Some growth retardants has an influence on the level and/or stability of the pH in the pot.
4. Some growth retardants can be mixed together with comparable results. F.i. Daminozide and Cycocel.

## GROWTH RETARDENTS (DAMINOZIDE)

1. Powder – 3 hours waiting after making the solution.
2. Granulate and liquid – can be used directly.
3. In a concentrated form, never more than 142 gram per liter.
4. Storage in cool and dark conditions (cool store).
5. Preference: No mixing with other chemicals.
6. Check the pressure (bar) before spraying, can cause residu on leaves.
7. Every treatment give a 3 - 4 day delay of the crop.
8. Crucial: mist treatment in Long Days, induction period in Short Days.
9. Spraying in the morning, most active period.
10. Moisty conditions better than dry conditions.
11. Give one hour after spraying a shower of water, this increases the effect.

## SCHEDULE FOR DAMINOZIDE TREATMENT (BASIC)

| SCHEDULE FOR DAY INCLUDE TREATMENT (BASIS) |   |   |              |   |   |   |   |                                    |   |   |   |    |
|--------------------------------------------|---|---|--------------|---|---|---|---|------------------------------------|---|---|---|----|
| Long days                                  |   |   | Short days   |   |   |   |   | Flowering                          |   |   |   |    |
| 1                                          | 2 | 3 | 1            | 2 | 3 | 4 | 5 | 6                                  | 7 | 8 | 9 | 10 |
| ▲                                          |   |   | if necessary |   |   |   |   | ▲                                  |   |   |   |    |
| Flowering misting treatments               |   |   |              |   |   |   |   | If necessary                       |   |   |   |    |
|                                            |   |   |              |   |   |   |   | Treatments during induction period |   |   |   |    |

## **SPRAYING METHODS**

1. Misting treatment – Very light, low volume treatment. Only the top of the plant is treated, not the lower levels.
2. Spray treatment – Standard treatment, volume to run-off. All levels of the plant are treated.
3. Irrigation treatment – High volume treatment, run-off. A lot of fluid also ends up in the pot.

## **LONG DAY PERIOD**

1. Slow growing varieties (Goldstrike): 0 - 1 x misting treatment.
2. Normal growing varieties (Mikeno/Rubio): 1 x misting treatments.
3. Strong growing varieties (Coto/Whites): 1 x - 2 x misting treatments.

## **SHORT DAY PERIOD (THE INDUCTION PERIOD)**

1. Varieties with a slow responstime (Goldstrike/Weaver): spray treatments in the 4<sup>th</sup> and 5<sup>th</sup> week.
2. Varieties with a normal responstime (Mikeno/Rubio): spray treatments in the 3<sup>rd</sup> and 4<sup>th</sup> week.
3. Varieties with a quick responstime (Coto/Birkin): spray treatments in the 3<sup>rd</sup> 4<sup>th</sup> and 5<sup>th</sup> week.

## **DIFFERENCE BETWEEN DAMINOZIDE & BONZI**

| DAMINOZIDE                                | BONZI                                 |
|-------------------------------------------|---------------------------------------|
| 1. Works mainly through leaves            | 1. Works mainly by roots              |
| 2. Starts working in 2 - 3 days           | 2. Starts working in 6 - 7 days       |
| 3. Works cumulative 3                     | 3. Works every time separate          |
| 4. Fixed                                  | 4. Rate depends on time of the year   |
| 5. Trustworthy for culture                | 5. Unreliable                         |
| 6. Delay culture 3 - 4 days per treatment | 6. Delay can be weeks with wrong rate |

## **CROP PROTECTION**

1. Determination of the problem/disease
2. No liquid chemicals, unless marked safe in cultural guides.
3. No dusty chemicals allowed.
4. With a scouting policy only curative treatments necessary.
5. Be very carefull with wetting agents.
6. No cocktails with growth retardents.
7. Avoid sprays on opening buds and flowering plants.
8. If a problem is treated, finish the schedule (especially with insects).
9. Most chemicals spray in the morning, except contact chemicals for insects, spray on the end of the day.

## **BIOLOGICAL CONTROL**

1. Determination of the problem/disease.
2. Scouting determines the action necessary.
3. Climatical conditions important for biological controllers.
4. Follow the schedule in detail: timing is important.
5. Sulphur evaporation not allowed

## **MOTHER STOCK CULTURE**

1. Requirements roughly comparable with Long Day culture finished plants.
2. Different ages of motherplants requires different feeding levels.
3. Assimilation lighting required to bring uniformity throughout the year (northern areas).
4. Early maintenance (e.g. pinching) of plants essential.
5. Maintenance throughout the culture (e.g. removing big leaves, misformed shoots etc) very important for production.
6. Scouting on diseased plants and plants which look different. If it looks different, remove it.
7. Harvesting by preference with a knife.
8. Harvesting cuttings always from the top of the plant – uniformity.
9. Harvesting cuttings on regular basis – at least once a week.
10. First three flushes – very heavy growing of cuttings.
11. Production after 7 months – cutting quality is decreasing.
12. The perfect cutting overall – 1 larger leafpair + 1 smaller leafpair + Top of the cutting.
13. Cutting too small (f.i. one bigger leafpair) is sensitive for fall-out, more difficult to root.
14. Cutting too big (f.i. with 3 large leafpairs) causing slower growth, less branching, sensitive for fall-out.
15. Selection checks and renewals by year.

## **DECORATIVE (CALANDIVA®) CULTURE**

1. Requirements roughly comparable with single flower culture.
2. Long Days do have the same periods.
3. Short Days must be at least 7 weeks.
4. Culture is in average 1,5 week longer than single flower culture.
5. The decoratives must have 14 hours of darkness for induction.
6. The decoratives can stand a higher light level.

## **OUTDOOR CULTURE (Calor; Deck`d Out)**

1. Requirements roughly comparable with single flower culture.
2. Suited for larger potsizes, 1 cutting in 13 cm – 17 cm pot (5-7 inch)
3. Long Days depending on pot size.
4. Short Days must be at least 7 weeks.
5. Culture is in average 1 week longer than single flower culture.
6. No pinching.
7. The Calor series are ideal for outdoor use, even in the full sun.
8. Hardly no growth-retardents required.

### **CULTURE IN HIGH TEMPERATURES (OVER 35 °C/90 °F)**

1. Try to cool down the plant at high noon with overhead water.
2. Keep the soil moisty (not wet), to avoid cupping.
3. Don't use cold water, below 10°C (50 °F), for this purpose.
4. Fertilization level in the pot not over 1.0 EC.
5. Try to keep some humidity in the greenhouse, at least 50%.
6. Avoid too much direct sunlight, shading is very important.
7. Start shading at 45.000 lux, to avoid cupping.
8. Open the blackout clothing if possible, watch out for interfering light.
9. Interfering light can also be the moon.
10. Spraying in the very early morning.
11. Act in time with growth-regulators, to avoid heat-stretching.

If you need advice, please contact **Ike Vlieland**  
**[i.vlieland@dummenorange.com](mailto:i.vlieland@dummenorange.com)**

**De Lier, 2021**

## How to install the app:



**For Android devices:**

- 1 Scan the QR code or visit our download location:  
[www.dummenorange.com/onTarget/getapp](http://www.dummenorange.com/onTarget/getapp)
- 2 Download and install our App
- 3 Enable your browser to allow the installation of unknown apps. Go to: Apps > Chrome > Install unknown apps > Allowed
- 4 Sign up for a free account
- 5 After approval you get full access to all features



**For Apple devices:**

- 1 Send an email to [DummenOrangeApp@DummenOrange.com](mailto:DummenOrangeApp@DummenOrange.com) mentioning that you wish you to use our app
- 2 You will receive an email from TestFlight with instructions and an install link
- 3 Follow the instructions to install our app on your device
- 4 Sign up for a free account
- 5 After approval you get full access to all features

**DummenOrange.com**

# GROWTH SUPPORT

from Dümme Orange

This cultivation tool supports growers to produce floricultural products in the best possible way.

It provides growth strategies, information about pests and diseases, a logbook for observations and measurements and a direct chat option with our technical experts.

This app helps you run your business with lower risks and higher future margins. The app also works offline and colleagues can collaborate in the logbook. Everyone can use this app to browse through our complete assortment. Our customers have exclusive access to all features.

for you

f DümmeOrangeNA  
f DümmeOrangeEU  
v Dümme Orange  
in Dümme Orange

Featuring:



QUICK START GUIDE



GROWTH STRATEGIES




COMPLETE ASSORTMENT



GROWTH CYCLE




PLANT PATHOLOGY



TECHNICAL SUPPORT




Add observations, notes and pictures to your Growth Cycle timeline



Identify pests and diseases with our easy-to-read expert knowledge and get immediate advice




Recommended growing conditions for your crop



Scan this QR-code to get this app on your Android device

QUICK START GUIDE

Containing the essential basics to grow this crop



MY QUESTIONS

Chat with our experts to get hands-on support




MY CATALOGUE

Browse, search and filter through our vast range of flowers and plants



MY TEAM

Collaborate with your team on your Growth Cycles



Download and print QR-codes to use in your greenhouse, so you can immediately access variety data with a simple scan