

Homerun Family

Chrysanthemum x morifolium



The Homerun family has perfectly matched colors that are well suited for monoculture or combinations. This family will work well for both shaded and natural season production. This family boasts very large flowers and strong color retention even under heat These mums have a medium to large vigor and have an expected natural response flowering window around week 38 (Late September).



65116
White



65378
Yellow



65911
Pink



65414
Scarlet



65113
Orange



48318
Confetti Garden Ruth



48319
Confetti Garden Mays



48317
Confetti Garden Robinson



48320
Confetti Garden Mantle

STATUS	CULTIVAR	CULTIVAR CODE	RESPONSE TIME (North/South/ Shaded)	VIGOR	FLOWER SIZE	HABIT	RECOMMENDED CONTAINER SIZE
TOP PICK	Pink	65911	37/38/6	M	L	M	Pint, Qt, 3Qt, Patio Pot, Hanging Basket
TOP PICK	Orange	65113	37/38/6	M	L	M	Pint, Qt, 3Qt, Patio Pot, Hanging Basket
	Scarlet	65414	38/38/6	M	L	M	Pint, Qt, 3Qt, Patio Pot, Hanging Basket
TOP PICK	White	65116	37/38/6	M	L	M	Pint, Qt, 3Qt, Patio Pot, Hanging Basket
TOP PICK	Yellow	65378	37/38/6	M	L	M	Pint, Qt, 3Qt, Patio Pot, Hanging Basket
	Confetti Garden Mantle	48320	37/38/6	M	L	M	3Qt. Patio Pot, Hanging Basket
	Confetti Garden Mays	48319	37/38/6	M	L	M	3Qt. Patio Pot, Hanging Basket
	Confetti Garden Robinson	48317	37/38/6	M	L	M	3Qt. Patio Pot, Hanging Basket
	Confetti Garden Ruth	48318	37/38/6	M	L	M	3Qt. Patio Pot, Hanging Basket

Homerun Family

Chrysanthemum x morifolium

STICKING

Rooting Hormone?	No
Misting instructions	Keep leaves coated and turgid (without saturating the media) until callus formation

CONTAINER SIZE	PLANTS PER POT	VEGETATIVE TIME (WEEKS)
4-6 in. (10-15 cm)	1	8
Gallon-8 in. (16-19 cm)	1	10
10 in. (22 cm) and up	3-5	10

FERTILITY

Fertilizer Rate	50 ppm N with 17-5-17 if media leaching occurs
Long-Day EC Range	Moderate (0.8-1.5 mS via SME)
pH Range	5.8-6.2

TEMPERATURE

Average Soil Temp	73-75°F (23-24°C)
Average Air Temp	70°F (21°C)

IRRIGATION

Irrigation Frequency (after day 14)	Level 3: Keep media moderately moist at all times
-------------------------------------	--

LIGHT

Vegetative Photoperiodic Requirements	Long Day
Supplemental	To prevent premature bud set, provide long days using a 4-hour night interruption from 10pm-2am at a minimum of 10 fc (108 lux)
Benefits From Shade?	Keep light levels below 4200 fc (45000 lux) until callus forms. Full sun conditions should persist thereafter

PINCHING

Not recommended

POTENTIAL DISEASES

Pythium, Fusarium, Botrytis, Rhizoctonia, Septoria, Alternaria, BLS - bacterial leafspot, white rust,

POTENTIAL PESTS

thrips, aphids, mealy bugs, mites, leafminers,

GROWTH REGULATION

- If nighttime temperatures fall below 68°F (20°C), apply Ethephon (3.9%) at 500 ppm once roots are about 1 inch (2.5 cm) in length (around day 10)

TIPS

- Place cuttings in cooler upon arrival at 46-50°F (8-10°C).

Homerun Family

Chrysanthemum x morifolium



FERTILITY

Fertilizer Rate	Moderate: 100-200 ppm N with 20-10-20
Short Day EC Range	Moderate-High (1.7-3.0 mS via SME)
pH Range	5.3 - 6.6

TEMPERATURE

Day Temperature	76-84°F (24-29°C)
Night Temperature	69-74°F (21-23°C)

IRRIGATION

Irrigation Frequency	Level 3: Keep media moderately moist at all times
----------------------	--

LIGHT

Flowering Photoperiodic Requirements	Short Day
Photoperiodic Instructions	Between 12.75-13.5 hours of darkness until flower color is visible.
Supplemental	Short days are required for bud set, only light to maintain vegetative growth
Benefits From Shade?	No

PINCH INSTRUCTIONS

Not recommended

POTENTIAL DISEASES

pythium, fusarium, botrytis, rhizoctonia, septoria, alternaria, bacterial leaf spot, white rust

POTENTIAL PESTS

thrips, aphids, mealy bugs, mites, leafminers

GROWTH REGULATION

- Ethephon (3.9%) at 500ppm every 7-10 days when temps are above 65°F (18°C); at 500-750ppm every 5-7 days when temps are between 55-65°F (12-18°C); and at 750-1000 ppm every 5-10 days when temps are below 54°F (11°C). All applications should stop a half week earlier than the response timing (e.g. last application 5.5 weeks before ship for a 5 week response time variety).
- GA4A7, 6-BA (1.8%, 1.8%) tank mix at 1:100 ratio with Ethephon (3.9%) to avoid brittle plants and an overly compact plant habit
- Daminozide (85%) at 500-1500 ppm for height control
- Paclobutrazol (0.4%) drench at 2-4ppm or spray at 30-125 ppm to hold at finish

TIPS

- Cool night temperatures during finishing can increase color intensity.
- Additional micronutrients from a balanced blend should be added to the fertilizer solution to deliver about 1 ppm Iron, 0.5 ppm Zinc, 0.25 ppm Copper, 0.25 ppm Boron, and 0.1 ppm Molybdenum.
- Heat delay can occur when daytime temperatures surpass 90°F (32°C) and/or when night temperatures surpass 77° (25°C). This is a general guideline and will differ based on cultivar and growing environment. In shaded production, ensure black cloth is pulled after sun is low to avoid heat buildup; evening or prior to sunrise is preferred.
- Irrigate with fertilizer immediately following a heavy rain to recharge the growing media. There is no reason to allow the media to dry before recharge.
- Plants not subjected to growth regulator will elongate about 1 inch (2.5 cm) per week following the start of short days. Plant height (including the height of the pot) should be about 60% of the final height at the start of short days and reproductive development.